



CRUCIFERS

CRUCIFERS (Cabbage, Cauliflower, Broccoli)


GROWTH STAGES	Seedbed		Trans-planting	Vegetative				Heading to Head Development			Maturity	
	1-15 DAS	16-25 DAS	1-7 DAT	8-14 DAT	15-21 DAT	22-28 DAT	29-35 DAT	36-42 DAT	43-49 DAT	50-56 DAT	57-70 DAT	
INSECTS	Cutworms / Armyworms											
	Diamond-back Moths											
	Webworms											
	Cabbage Worms											
	Leafminers											
RECOMMENDED INSECTICIDES USE RATE / 16L water tank load	SUMICIDIN 5 - 8 Tbsp		Padan 1.5 - 2 Tbsp		SUCCESS[®] NATURALYTE 2.5 - 5.0 Tbsp		Padan 1.5 - 2 Tbsp		SUCCESS[®] NATURALYTE 2.5 - 5.0 Tbsp		Padan 1.5 - 2 Tbsp	SUMICIDIN 5 - 8 Tbsp
DISEASES	Seedling Diseases						Downy Mildew					
							Alternaria Leaf Spot					
RECOMMENDED FUNGICIDES USE RATE / 16L water tank load	4-8 Tbsp CAPTAN 50 WP		3-6 Tbsp				VONDOZEB PLUS					

INSECTS



CUTWORMS ARMYWORMS DIAMOND-BACK MOTHS WEBWORMS CABBAGE WORMS LEAFMINERS

DISEASES



DAMPING-OFF DOWNY MILDEW ALTERNARIA LEAF SPOT

SPECIALIST PRODUCTS USE RATE / 16L WATER TANK LOAD				
SPECIALIZATION	FOR PRE-PLANT WEED CLEARING	FOR CONTROL OF SLUGS	FOR FARMERS WHO PRACTICE INSECTICIDE TANK-MIXING	FOR LEAF SPOT
RECOMMENDED SPECIALIST	SLASH 16-36* Tbsp	BAYONET	SUMICIDIN 5-8 Tbsp	SAPROL 3-8 Tbsp

